

TAG-FORTE

Topical Anesthetic Dental Gel – Pre Injection

TAG-FORTE is a Dentist's only topical anaesthetic that contains the highest concentration of 25% Lidocaine ensuring rapid onset of active pain relief whilst maximising duration of action.

Indicated as a *topical anaesthetic* for the relief of pain associated with dental procedures within the oral cavity. TAG-FORTE is applied to the site of injection 1 - 2 mins before a procedure begins to relieve pain.

Our proprietary base utilises micelle technology to ensure penetration of lidocaine for improved dose delivery. Micelles encapsulate the lidocaine molecules and pass them across the mucosal epithelial layer (diagram 1.0).




Diagram 1.0

Efficient delivery of lidocaine across epithelial tissue will yield a faster onset of action giving your practice reduced wait time. The dosage of lidocaine is also improved leading to more effective pain relief.

TAG

Topical Anesthetic Dental Gel – Pre Injection

TAG is a Dentist's only topical anaesthetic that contains the highest concentration of 9.8% Lidocaine ensuring rapid onset of active pain relief whilst maximising duration of action.

Indicated as a *topical anaesthetic* for the relief of pain associated with dental procedures within the oral cavity. TAG is applied to the site of injection 1 - 2 mins before a procedure begins to relieve pain.

Our proprietary base utilises micelle technology to ensure penetration of lidocaine for improved dose delivery. Micelles encapsulate the lidocaine molecules and pass them across the mucosal epithelial layer (diagram 1.0).




Diagram 1.0

Efficient delivery of lidocaine across epithelial tissue will yield a faster onset of action giving your practice reduced wait time. The dosage of lidocaine is also improved leading to more effective pain relief.

TAG-FORTE Topical Anesthetic Dental Gel

MATERIAL SAFETY DATA SHEET

Trade Name:	TAG-FORTE
Chemical Name:	Lidocaine 25%
Product Definition:	Indicated as a topical anaesthetic prior to dental injection(s) for the relief of pain associated with dental procedures within the oral cavity Topical application for the extensive desensitisation of gingivi prior to dental hygiene procedures
Supplier's Details:	Como Compounding Pharmacy 299 Toorak Rd South Yarra VIC 3141 PH: 1300 889 173
Contact:	Nima Alavi B.Pharm

PHYSICAL DATA

Appearance and Odour: Cloudy colourless gel, with peppermint flavouring

SUPPLIER DISCLAIMER - FOR DENTAL USE ONLY

Use as directed. The information and recommendations related to Lidocaine have been taken from MIMS 2011.

TAG-FORTE Topical Anesthetic Dental Gel

MATERIAL SAFETY DATA SHEET

Usage Instructions:

TAG-FORTE is to be applied **sparingly** to the site of injection and onset can be expected within 1 -2 mins.

TAG-FORTE is contraindicated in patients with a history of sensitivity to local anaesthetics (of the amide type).

Storage & Disposal

Keep in cool, dry environment. Store & dispose of TAG-FORTE out of the reach of children, pets and others.

HEALTH HAZARD DATA

Excessive Dosing & Toxicity

Excessive dosing i.e. applying TAG-FORTE to larger areas than the injection site could result in absorption of lidocaine into the blood. High blood levels of lidocaine can be toxic. Local irritation to TAG-FORTE could be due to lidocaine or could be attributed to one of the inactive ingredients.

Emergency & First Aid Procedures

In case of suspected toxicity or hypersensitivity call Poisons Info on 131126 or contact a physician.

Eyes

Avoid contact of TAG-FORTE with the eye. Will cause eye irritation and reddening of the eye.

Emergency & First Aid Procedures

In case of eye contact, flush immediately and thoroughly for a minimum of 15 mins with water or saline and seek medical advice.

TAG Topical Anesthetic Dental Gel

MATERIAL SAFETY DATA SHEET

Trade Name:	TAG
Chemical Name:	Lidocaine 9.8%
Product Definition:	Indicated as a topical anaesthetic prior to dental injection(s) for the relief of pain associated with dental procedures within the oral cavity Topical application for the extensive desensitisation of gingivi prior to dental hygiene procedures
Supplier's Details:	Como Compounding Pharmacy 299 Toorak Rd South Yarra VIC 3141 PH: 1300 889 173
Contact:	Nima Alavi B.Pharm

PHYSICAL DATA

Appearance and Odour: Cloudy colourless gel, with tutti-frutti flavouring

SUPPLIER DISCLAIMER - FOR DENTAL USE ONLY

Use as directed. The information and recommendations related to Lidocaine have been taken from MIMS 2011.

TAG Topical Anesthetic Dental Gel

MATERIAL SAFETY DATA SHEET

Usage Instructions:

TAG is to be applied **sparingly** to the site of injection and onset can be expected within 1 -2 mins.

TAG is contraindicated in patients with a history of sensitivity to local anaesthetics (of the amide type).

Storage & Disposal

Keep in cool, dry environment. Store & dispose of TAG out of the reach of children, pets and others.

HEALTH HAZARD DATA

Excessive Dosing & Toxicity

Excessive dosing i.e. applying TAG to larger areas than the injection site could result in absorption of lidocaine into the blood. High blood levels of lidocaine can be toxic. Local irritation to TAG could be due to lidocaine or could be attributed to one of the inactive ingredients.

Emergency & First Aid Procedures

In case of suspected toxicity or hypersensitivity call Poisons Info on 131126 or contact a physician.

Eyes

Avoid contact of TAG with the eye. Will cause eye irritation and reddening of the eye.

Emergency & First Aid Procedures

In case of eye contact, flush immediately and thoroughly for a minimum of 15 mins with water or saline and seek medical advice.
